

Answer to Dermacase *continued from page 247*

4. Favre-Racouchot syndrome

Favre-Racouchot syndrome (FRS), first noted by Favre in 1932,¹ and detailed by Favre and Racouchot in 1951,² is a condition characterized by cysts, comedones, and nodular elastosis in sun-damaged skin. It most commonly presents in elderly white men with a history of long-term sun exposure and heavy smoking.^{3,4} Favre-Racouchot syndrome is a disease of cosmetic concern, limited to the skin with no related internal manifestations.

Epidemiology

Favre-Racouchot syndrome is not an uncommon condition—it occurs in up to 6% of those older than 50 years of age.⁵ The incidence of FRS increases with age, although it has been reported in patients as early as in their second decade of life.⁶ Prevalence is highest in white men, but the condition has been observed in women and dark-skinned individuals.^{7,8} Although the exact mechanism of the condition is not known, FRS has been specifically connected to sun exposure,⁹ smoking,³ and, in a minority of cases, radiation exposure.¹⁰

Etiology and pathogenesis

Favre-Racouchot syndrome is characterized by solar elastosis, with the presence of nodules, cysts, and comedones. Solar elastosis refers to the damage to dermal elastic tissue due to prolonged exposure to UVA and UVB rays. The histologic changes present in FRS include an atrophic epidermis and large masses of keratinous material, causing follicular plugging, as well as solar elastotic changes (which include increased elastic tissue with thickened, tortuous fibres in the upper and mid dermis) and possible basophilic degeneration of connective tissue.⁸ Although it is suggested that senile comedones and sebaceous gland hyperplasia are somehow associated with photodamage, no proof of this has been documented.¹¹ Interestingly, the comedones found in FRS are histologically indistinguishable from the primary comedones of acne vulgaris, with the exception of a lack of inflammation and the presence of a marked actinic elastosis in the surrounding dermis.^{10,12} It is hypothesized that damage to the dermal elastic network and subsequent reduction of tensile strength might be what predisposes these comedones to histologic change, expansion, and subsequent formation of cysts and nodules.^{6,8}

Diagnosis and differential


Favre-Racouchot syndrome can be diagnosed by clinical findings of multiple open and closed comedones with yellowish nodules of elastotic material in patients older than 40 years of age. Patients with FRS usually present with a history of long-term sun exposure and smoking, and sometimes the possibility of radiation exposure, although this is rare. Clinically, examination

reveals histologic changes manifested as actinically damaged skin with atrophy; yellowish discolouration; wrinkles and furrows; cystic nodules; and punctate, waxy, noninflamed, soft, open or closed comedones.^{4,8} As it is hypothesized that the underlying etiology of FRS is UV damage, lesions will generally be photodistributed. Affected areas of the body are primarily the periorbital and temporal areas, but the malar eminences, lateral neck, earlobes, and postauricular areas might also be involved.⁴ The eruptions are usually bilateral and symmetric but can be asymmetric or unilateral, depending on the history of sun exposure.¹³ As a member of the spectrum of skin conditions caused by long-term sun exposure known as dermatoheliosis, FRS can be associated with actinic keratoses, basal and squamous cell carcinoma, cutis rhomboidalis nuchae, trichostasis spinulosa, and keratoacanthoma.

The differential diagnosis for FRS includes dermatoses that present with lesions of similar morphologies, but it can nonetheless be distinguished based on a variety of features. These features include the following: acne vulgaris (lesions limited to inflamed comedones); sebaceous hyperplasia (lesions limited to enlarged sebaceous glands); actinic comedonal plaques (similar lesions, also found on the flexural surface of the arms and often unilateral); cutis rhomboidalis nuchae (similar skin morphology, differentiated by the presence of deep, crisscross furrows on the dorsal and lateral skin of the neck); chloracne (similar lesions, but also localized in postauricular areas, axillae, and groin); actinic granuloma (different lesional morphology: pink or skin-coloured papules in annular or serpiginous plaques with atrophic centres); colloid milium (different lesional morphology: small, amber, waxy, translucent, and firm papules occurring in groups); and syringomas or trichoepitheliomas (both presenting with similar lesions but occurring in a younger patient population compared with FRS).^{4,14}

Treatment

It is important to note that, unless treated, FRS is slowly progressive. First, it is key that all patients are advised to take proper sun precautions, such as wearing sunscreen with a sun protective factor of at least 30 with UVA and UVB protection and avoiding the outdoors between 10 AM and 2 PM if at all possible. If the patient is a smoker, quitting is strongly advised. In treating the patient, 2 different approaches can be used: pharmacologic and surgical. Topical retinoids, such as tretinoin, adapalene, or tazarotene, are the most effective pharmacologic treatments. Tretinoin, in addition to eliminating comedones, has been shown to reduce the aged appearance of sun-damaged skin and has beneficial effects on collagen.¹⁵ Retinaldehyde, a precursor to retinoic acid, shows similar effectiveness; in addition, it has been shown to increase the surface area of elastin and

collagen fibres in UVA-exposed skin.¹⁶ Surgical techniques include excision, dermabrasion, curettage, comedone extraction, and laser resurfacing. Although these techniques have yielded poor results when used independently, they allow FRS a very favourable outcome when used in conjunction with medication.^{4,14} 

Mr Sawicki is a third-year medical student at Queen's University in Kingston, Ont. Dr Barankin is a dermatologist practising in Toronto, Ont.

Competing interests

None declared

References

1. Favre M. Sur une affection kystique des appareils pilosébacés localisée à certaines régions de la face. *Bull Soc Fr Dermatol Syphiligr* 1932;39:93-6.
2. Favre M, Racouchot J. Nodular cutaneous elastoidosis with cysts and comedones [article in French]. *Ann Dermatol Syphiligr* (Paris) 1951;78(6):681-702.
3. Keough GC, Laws RA, Elston DM. Favre-Racouchot syndrome: a case for smokers' comedones. *Arch Dermatol* 1997;133(6):796-7.
4. Patterson WM, Fox MD, Schwartz RA. Favre-Racouchot disease. *Int J Dermatol* 2004;43(3):167-9.
5. Friedman SJ, Su WP. Favre-Racouchot syndrome associated with radiation therapy. *Cutis* 1983;31(3):306-10.
6. Jansen T, Plewig G. Favre-Racouchot disease. In: Demis DJ, editor. *Clinical dermatology*. Philadelphia, PA: Lippincott Williams and Wilkins; 1998. p. 4-44.
7. Schäfer T, Merkl J, Klemm E, Wichmann HE, Ring J; KORA Study Group. The epidemiology of nevi and signs of skin aging in the adult general population: results of the KORA-survey 2000. *J Invest Dermatol* 2006;126(7):1490-6. Epub 2006 Apr 27.
8. Lewis KG, Bercovitch L, Dill SW, Robinson-Bostom L. Acquired disorders of elastic tissue: part I. Increased elastic tissue and solar elastotic syndromes. *J Am Acad Dermatol* 2004;51(1):1-21; quiz 22-4.
9. Heras JA, Jimenez F, Soguero ML, Arjona C, Plaza JA, Martinez P, et al. Bullous solar elastosis. *Clin Exp Dermatol* 2007;32(3):272-4. Epub 2007 Jan 23.
10. Breit S, Flaig MJ, Wolff H, Plewig G. Favre-Racouchot-like disease after radiation therapy. *J Am Acad Dermatol* 2003;49(1):117-9.
11. Kumar P, Marks R. Sebaceous gland hyperplasia and senile comedones: a prevalence study in elderly hospitalized patients. *Br J Dermatol* 1987;117(2):231-6.
12. Sanchez-Yus E, del Rio E, Simon P, Requena L, Vazquez H. The histopathology of closed and open comedones of Favre-Racouchot disease. *Arch Dermatol* 1997;133(6):743-5.
13. Mavilia L, Rossi R, Cannarozzo G, Massi D, Cappugi P, Campolmi P. Unilateral nodular elastosis with cysts and comedones (Favre-Racouchot syndrome): report of two cases treated with a new combined therapeutic approach. *Dermatology* 2002;204(3):251.
14. Cuce LC, Paschoal LH, Curban GV. Cutaneous nodular elastoidosis with cysts and comedones. *Arch Dermatol* 1964;89:798-802.
15. Kligman AM, Plewig G, Mills OH Jr. Topically applied tretinoin for senile (solar) comedones. *Arch Dermatol* 1971;104(4):420-1.
16. Fluhr JW, Vienne MP, Lauze C, Dupuy P, Gehring W, Gloor M. Tolerance profile of retinol, retinaldehyde and retinoic acid under maximized and long-term clinical conditions. *Dermatology* 1999;199(Suppl 1):57-60.

— * * * —